



Cortical Neuroplasticity and Cognition in Extreme Environments

Anika Friedl-Werner

Charité - Universitätsmedizin Berlin, Institute of Physiology, Center for Space Medicine and Extreme Environments Berlin, Berlin, Germany & Université de Caen Normandie, COMETE, INSERM U 1075, Caen, France

Abstract

Space is one of the most extreme environments for human beings, yet space faring nations strive for deep space exploration and human settlement on Moon and Mars. These missions expose crews to various physiological and psychological stressors such as radiation, altered gravity conditions, social isolation and confinement, and reduced physical activity - stressors that are also known to alter neuroplasticity and cognitive functions. To ensure missions' success and safety, it is important to know how the brain adapts in response to these stressors and to understand the associated neurobehavioral risks.

As part of this dissertation, the impact of three of those stressors on brain function was examined through two different spaceflight analog models, i.e., *Head-down tilt Bed Rest* and *Parabolic Flight*. The effects of *Physical Inactivity* in form of long-term bed rest on episodic memory and its neural correlates by means of functional magnetic resonance imaging were the focus of *Research Paper I*. The impact of long-term head-down tilt bed rest, which is also characterized as a *semi-isolated, confined and sensory deprived environment*, on affective processing and its underlying electrocortical activity was investigated in *Research Paper II*. In *Research Paper III*, the impact of short bouts of *Microgravity* on attentional processes during a parabolic flight was examined.

First, the results showed that long-term immobilization of two months of Head-down tilt Bed Rest (HDBR) altered brain activity during memory encoding and retrieval in the left

hippocampus and parahippocampal gyrus. These changes could be largely counteracted by a high-intensity training intervention that was performed during the immobilization period. Second, bed rest associated social isolation and confinement evoked reduced Event Related Potential (ERP) amplitudes in participants while looking at highly arousing photographs. This emotional blunting was observed predominantly in centroparietal regions and did not occur in a control group. Source localization confirmed a lower electrocortical activity in the posterior cingulate gyrus, insula, and precuneus in the bed rest group for pleasant and unpleasant, but not for neutral photographs. Third, attentional performance during a Continuous Performance Task (CPT) was impaired by weightlessness. There was additional evidence that attentional performance was also influenced by participants' emotional states.

The results provide evidence of the adverse neurobehavioral adaptations brought about by these spaceflight associated stressors. The results go beyond applications of space medicine and provide further insight into the adaptational processes of the brain in response to physical inactivity, confinement, and vestibular deficiency.